



Generalized variational principles for micromorphic magnetoelectroelastodynamics

Cheng-Bo Zheng, Bin Liu, Zuo-Jun Wang^{*}, Hong-Shi Lü

Key Laboratory of Measurement Technology and Instrumentation of Hebei Province, Yanshan University, Qinhuangdao 066004, PR China

ARTICLE INFO

Keywords:

Micromorphic
magnetoelectroelastodynamics
Generalized variational principle
Semi-inverse method

ABSTRACT

A family of generalized variational principles is established for the initial-boundary value problem of micromorphic magnetoelectroelastodynamics by He's semi-inverse method. This paper aims at providing a more complete theoretical basis for the finite element applications.

© 2010 Elsevier Ltd. All rights reserved.

1. Introduction

Magnetoelectroelastic materials are gaining attention because of their capability of converting energies among the magnetic, electric and mechanical forms. Many variational principles have been established in piezoelectricity [1–5], thermopiezoelectricity [6], piezoelectromagnetism [7,8], and magnetoelectroelasticity [9–13]. Recently Nappa [14] and He [15,16] established the variational principles in micromorphic thermoelasticity [17].

In this paper, on the basis of He's contributions [9,16], we will deduce a family of generalized variational principles for the initial-boundary value problem of micromorphic magnetoelectroelastodynamics by He's semi-inverse method [18–21].

2. Fundamental equations

Let Ω be a 3D regular region of an elastic continuum with the piecewise smooth surface $\partial\Omega$. The fundamental equations for the micromorphic behavior [17,14–16] of elastic bodies, coupled with quasi-static electromagnetic fields, consist of the equations of motion

$$\sigma_{;j}^i + F^i = \dot{p}^i, \quad (1)$$

$$m_{;k}^{kij} + \sigma^{ji} - s^{ji} + L^{ij} = \dot{\pi}^{ij}, \quad (2)$$

$$p^i = \rho v^i, \quad (3)$$

$$\pi^{ij} = \rho l_k^j \varpi^{ik}, \quad (4)$$

$$v_i = \dot{u}_i, \quad (5)$$

$$\varpi_{ij} = \dot{\phi}_{ij}, \quad (6)$$

the constitutive equations

$$\sigma^{ij} = C_{\varepsilon\varepsilon}^{ijmn} \varepsilon_{mn} + C_{\varepsilon e}^{ijmn} e_{mn} + C_{\varepsilon\gamma}^{ijmnp} \gamma_{mnp} - C_{\varepsilon E}^{mij} E_m - C_{\varepsilon B}^{mij} B_m, \quad (7)$$

^{*} Corresponding author.

E-mail address: scoutqhd@foxmail.com (Z.-J. Wang).

$$s^{ij} = C_{\varepsilon\varepsilon}^{ijmn} \varepsilon_{mn} + C_{ee}^{ijmn} e_{mn} + C_{e\gamma}^{ijmnp} \gamma_{mnp} - C_{eE}^{mij} E_m - C_{eB}^{mij} B_m, \quad (8)$$

$$m^{ijk} = C_{\varepsilon\gamma}^{mnijk} \varepsilon_{mn} + C_{e\gamma}^{mnijk} e_{mn} + C_{\gamma\gamma}^{ijkmp} \gamma_{mnp} - C_{\gamma E}^{mijk} E_m - C_{\gamma B}^{mijk} B_m, \quad (9)$$

$$D^i = C_{\varepsilon E}^{imn} \varepsilon_{mn} + C_{eE}^{imn} e_{mn} + C_{\gamma E}^{imnp} \gamma_{mnp} + C_{EE}^{im} E_m + C_{EB}^{im} B_m, \quad (10)$$

$$H^i = C_{\varepsilon B}^{imn} \varepsilon_{mn} + C_{eB}^{imn} e_{mn} + C_{\gamma B}^{imnp} \gamma_{mnp} + C_{EB}^{im} E_m + C_{BB}^{im} B_m, \quad (11)$$

the geometrical equations

$$\varepsilon_{ij} = u_{j;i} - \varphi_{ji}, \quad (12)$$

$$2e_{ij} = \varphi_{ij} + \varphi_{ji}, \quad (13)$$

$$\gamma_{ijk} = \varphi_{ij;k}, \quad (14)$$

the equations for electromagnetic fields

$$\varepsilon^{ijk} H_{k;j} = J^i, \quad (15)$$

$$D^i_{;i} = q, \quad (16)$$

$$B_i = \varepsilon_{ijk} A^{k;j}, \quad (17)$$

$$E_i = -\phi_{,i}, \quad (18)$$

the boundary conditions

$$u_i = \bar{u}_i \quad \text{on } \partial\Omega_u, \quad (19)$$

$$\sigma^{ij} n_j = \bar{T}^i \quad \text{on } \partial\Omega_\sigma, \quad (20)$$

$$\varphi_{ij} = \bar{\varphi}_{ij} \quad \text{on } \partial\Omega_\varphi, \quad (21)$$

$$m^{kij} n_k = \bar{M}^{ij} \quad \text{on } \partial\Omega_m, \quad (22)$$

$$n_i D^i = \bar{d} \quad \text{on } \partial\Omega_D, \quad (23)$$

$$\phi = \bar{\phi} \quad \text{on } \partial\Omega_\phi, \quad (24)$$

$$\varepsilon^{ijk} n_j H_k = \bar{h}^i \quad \text{on } \partial\Omega_H, \quad (25)$$

$$A_k = \bar{A}_k \quad \text{on } \partial\Omega_A, \quad (26)$$

and the initial conditions

$$u_{i0} = \bar{u}_{i0}, \quad (27)$$

$$p^{i0} = \bar{p}^{i0}, \quad (28)$$

$$\varphi_{ij0} = \bar{\varphi}_{ij0}, \quad (29)$$

$$\pi^{ij0} = \bar{\pi}^{ij0}. \quad (30)$$

Here σ^{ij} is the stress tensor, s_{ij} is the microstress tensor, m_{kij} is the stress moment tensor, u_i is the displacement vector, v_i is the velocity vector, p^i is the momentum vector, φ_{ij} is the microdeformation tensor, π^{ij} is the micromomentum tensor, ϖ^{ik} is the microvelocity tensor, F_i is the body force vector, L_{ij} is the body moment tensor, ε_{ij} , e_{ij} and γ_{ijk} are the linear strain tensors, ρ is the reference mass density, I_k^i is the microinertia, D^i is the electric displacement vector, B_i is the magnetic induction vector, E_i is the electric field intensity vector, H^i is the magnetic field intensity vector, ϕ is the scalar potential, A_i is the vector potential, q is the electric charge density, J^i is the electric current density vector, $C_{XY}^{ij\dots}$ are the constitutive coefficients, ε^{ijk} is the alternating tensor, ‘;’ denotes covariant differentiation.

3. Variational formulations

Using He’s semi-inverse method [18–21], we construct a trial functional in the form

$$J = \int_{t_0}^{t_1} dt \iiint_{\Omega} L dV + I_B, \quad (31)$$

where I_B is the integral boundary, and L is a trial Lagrangian, which is defined as

$$L = \frac{1}{2} \rho v^i v_i + \frac{1}{2} \rho I_k^j \varpi^{ik} \varpi_{ij} - \frac{1}{2} \varepsilon_{ij} C_{\varepsilon\varepsilon}^{ijmn} \varepsilon_{mn} - \frac{1}{2} e_{ij} C_{ee}^{ijmn} e_{mn} - \frac{1}{2} \gamma_{ijk} C_{\gamma\gamma}^{ijkmp} \gamma_{mnp} + \frac{1}{2} E_i C_{EE}^{im} E_m + \frac{1}{2} B_i C_{BB}^{im} B_m + f. \quad (32)$$

Here, $p^i, \pi^{ij}, \sigma^{ij}, s^{ij}, m^{kij}, D^i, H^i, v_i, \varepsilon_{ij}, \varpi_{ij}, \gamma_{ijk}, e_{ij}, B_i, E_i, u_i, \varphi_{ij}, A^i$ and ϕ are considered as independent fields, f is an unknown function. There exist alternative approaches for constructing the trial functional [22–26]. Let us introduce an internal energy density A , which is defined as

$$\begin{aligned} A(\varepsilon_{ij}, e_{ij}, \gamma_{ijk}, E_i, B_i) = & \frac{1}{2} \varepsilon_{ij} C_{\varepsilon\varepsilon}^{ijmn} \varepsilon_{mn} + \frac{1}{2} e_{ij} C_{ee}^{ijmn} e_{mn} + \frac{1}{2} \gamma_{ijk} C_{\gamma\gamma}^{ijk mnp} \gamma_{mnp} - \frac{1}{2} E_i C_{EE}^{im} E_m \\ & - \frac{1}{2} B_i C_{BB}^{im} B_m + \varepsilon_{ij} C_{\varepsilon e}^{ijmn} e_{mn} + \varepsilon_{ij} C_{\varepsilon\gamma}^{ijmnp} \gamma_{mnp} + e_{ij} C_{e\gamma}^{ijmnp} \gamma_{mnp} - \varepsilon_{ij} C_{\varepsilon E}^{mij} E_m - \varepsilon_{ij} C_{\varepsilon B}^{mij} B_m \\ & - e_{ij} C_{eE}^{mij} E_m - e_{ij} C_{eB}^{mij} B_m - \gamma_{ijk} C_{\gamma E}^{mijk} E_m - \gamma_{ijk} C_{\gamma B}^{mijk} B_m - E_i C_{EB}^{im} B_m. \end{aligned} \quad (33)$$

The Hu–Washizu-like Lagrangian can be written in the form

$$\begin{aligned} L_{HW} = & K - A - p^i(v_i - \dot{u}_i) - \pi^{ij}(\varpi_{ij} - \dot{\varphi}_{ij}) + F^i u_i + L^{ij} \varphi_{ij} + \sigma^{ij}(\varepsilon_{ij} - u_{i,j} + \varphi_{ji}) \\ & + s^{ij}(e_{ij} - \varphi_{ij}) + m^{kij}(\gamma_{ijk} - \varphi_{ij,k}) - D^i(E_i + \phi_{,i}) - H^i(B_i - \varepsilon_{ijk} A^{k,j}) - q\phi - J^i A_i, \end{aligned} \quad (34)$$

where K is the kinetic energy density, which is defined as

$$K = \frac{1}{2} \rho v^i v_i + \frac{1}{2} \rho I_k^j \varpi^{ik} \varpi_{ij}. \quad (35)$$

After incorporating the boundary conditions into the functional, we obtain a determined Hu–Washizu-like action, which reads

$$\begin{aligned} J_{HW} = & \int_{t_0}^{t_1} dt \iiint_{\Omega} L_{HW} dV + \int_{t_0}^{t_1} dt \iint_{\partial\Omega_u} (u_i - \bar{u}_i) \sigma^{ji} n_j dS + \int_{t_0}^{t_1} dt \iint_{\partial\Omega_m} \bar{M}^{ij} \varphi_{ij} dS \\ & + \int_{t_0}^{t_1} dt \iint_{\partial\Omega_\varphi} (\varphi_{ij} - \bar{\varphi}_{ij}) m^{kij} n_k dS + \int_{t_0}^{t_1} dt \iint_{\partial\Omega_\sigma} \bar{T}^i u_i dS + \int_{t_0}^{t_1} dt \iint_{\partial\Omega_\phi} (\phi - \bar{\phi}) D^i n_i dS \\ & + \int_{t_0}^{t_1} dt \iint_{\partial\Omega_D} \bar{d} \phi dS - \int_{t_0}^{t_1} dt \iint_{\partial\Omega_A} H^i \varepsilon_{ijk} n^j (A^k - \bar{A}^k) dS + \int_{t_0}^{t_1} dt \iint_{\partial\Omega_H} \bar{h}^i A_i dS \\ & + \iiint_V [-\hat{p}^{i1} u_{i1} + p^{i0} (u_{i0} - \bar{u}_{i0}) + (\bar{p}^{i0} - \hat{p}^{i0}) u_{i1} - \hat{\pi}^{ij1} \varphi_{ij1} + \pi^{ij0} (\varphi_{ij0} - \bar{\varphi}_{ij0}) + (\bar{\pi}^{ij0} - \hat{\pi}^{ij0}) \varphi_{ij1}] dV, \end{aligned} \quad (36)$$

where ‘ \wedge ’ denotes the restricted variation [27]. The action J_{HW} can directly lead to all the fundamental equations (1)–(30).

We can also introduce a complementary internal energy density B , which is defined as

$$\begin{aligned} B(\sigma_{ij}, s_{ij}, m_{ijk}, D_i, H_i) = & \frac{1}{2} \sigma_{ij} C_{\sigma\sigma}^{ijmn} \sigma_{mn} + \frac{1}{2} s_{ij} C_{ss}^{ijmn} s_{mn} + \frac{1}{2} m_{ijk} C_{mm}^{ijk mnp} m_{mnp} - \frac{1}{2} D_i C_{DD}^{im} D_m - \frac{1}{2} H_i C_{HH}^{im} H_m \\ & - \sigma_{ij} C_{\sigma s}^{ijmn} s_{mn} - \sigma_{ij} C_{\sigma m}^{ijmnp} m_{mnp} - s_{ij} C_{sm}^{ijmnp} m_{mnp} + \sigma_{ij} C_{\sigma D}^{mij} D_m + \sigma_{ij} C_{\sigma H}^{mij} H_m \\ & + s_{ij} C_{sD}^{mij} D_m + s_{ij} C_{sH}^{mij} H_m + m_{ijk} C_{mD}^{mijk} D_m + m_{ijk} C_{mH}^{mijk} H_m + D_i C_{DH}^{im} H_m. \end{aligned} \quad (37)$$

The Hellinger–Reissner-like Lagrangian can be written in the form

$$\begin{aligned} L_{HR} = & B - K(p^i, \pi^{ik}) + p^i \dot{u}_i + \pi^{ij} \dot{\varphi}_{ij} + F^i u_i + L^{ij} \varphi_{ij} \\ & + \sigma^{ij}(-u_{i,j} + \varphi_{ji}) - s^{ij} \varphi_{ij} - m^{kij} \varphi_{ij,k} - D^i \phi_{,i} + H^i \varepsilon_{ijk} A^{k,j} - q\phi - J^i A_i. \end{aligned} \quad (38)$$

Various variational principles can be deduced from the above obtained functionals. Here we give some Lagrangians for reference:

$$L_0 = K - A + F^i u_i + L^{ij} \varphi_{ij} - q\phi - J^i A_i, \quad (39)$$

$$L_1 = K - A - p^i v_i - \pi^{ij} \varpi_{ij} + s^{ij} e_{ij} + \sigma^{ij} \varepsilon_{ij} + m^{kij} \gamma_{ijk} - D^i E_i - H^i B_i, \quad (40)$$

$$\begin{aligned} L_2 = & K - A + (F^i - \dot{p}^i + \sigma_{,j}^{ij}) u_i - p^i v_i - \pi^{ij} \varpi_{ij} + (\sigma^{ij} - s^{ij} + m_{,k}^{kij} + L^{ij} - \dot{\pi}^{ij}) \varphi_{ij} \\ & + s^{ij} e_{ij} + \sigma^{ij} \varepsilon_{ij} + m^{kij} \gamma_{ijk} + (D_{,i}^i - q) \phi - D^i E_i - H^i B_i - (\varepsilon^{ijk} H_{k,j} + J^i) A_i. \end{aligned} \quad (41)$$

4. Conclusion

A family of generalized variational formations has been proposed for micromorphic magnetoelectroelastodynamics by He’s semi-inverse method. The present variational principles will provide a theoretical foundation for numerical techniques for the problem discussed.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (No. 60304009) and the Natural Science Foundation of Hebei Province, China (No. F2005000385).

References

- [1] R.A. Toupin, The elastic dielectrics, *J. Ration. Mech. Anal.* 5 (1956) 849–915.
- [2] M.C. Dökmeci, Variational principles in piezoelectricity, *Lett. Nuovo Cimento* 7 (1973) 449–454.
- [3] I.A. Vekovichsheva, Variational principles in the theory of electroelasticity, *Soviet Appl. Mech.* 7 (1974) 1049–1052.
- [4] J.H. He, Coupled variational principles of piezoelectricity, *Int. J. Eng. Sci.* 39 (2001) 323–341.
- [5] J.H. He, Minimum energy principles and generalized variational principles of nonlinear piezoelectricity, *J. Theor. Appl. Mech.* 31 (2) (2001) 89–96.
- [6] J.H. He, Hamilton principle and generalized variational principles of linear thermopiezoelectricity, *ASME J. Appl. Mech.* 68 (4) (2001) 666–667.
- [7] G. Altay, M.C. Dökmeci, Fundamental equations of certain electromagnetic acoustic discontinuous fields in variational form, *Contin. Mech. Thermodyn.* 16 (2004) 53–71.
- [8] G. Altay, M.C. Dökmeci, Variational principles and vibrations of a functionally graded plate, *Comput. & Structures* 83 (2005) 1340–1354.
- [9] J.H. He, Variational theory for linear magneto-electro-elasticity, *Int. J. Nonlinear Sci. Numer.* 2 (4) (2001) 309–316.
- [10] E. Luo, H.J. Zhu, L. Yuan, Unconventional Hamilton-type variational principles for electromagnetic elastodynamics, *Sci. China Ser. G* 49 (1) (2006) 119–128.
- [11] C.B. Zheng, B. Liu, Z.J. Wang, et al., Variational principle for nonlinear magneto-electro-elastodynamics with finite displacement by He's semi-inverse method, *Int. J. Nonlinear Sci. Numer.* 10 (2009) 1523–1526.
- [12] G. Altay, M.C. Dökmeci, On the fundamental equations of electromagnetoelastic media in variational form with an application to shell/laminae equations, *Internat. J. Solids Structures* 47 (2010) 466–492.
- [13] Z.J. Wang, D.Z. Zheng, C.B. Zheng, Simplified Gurtin-type generalized variational principles for fully dynamic magneto-electro-elasticity with geometrical nonlinearity, *Internat. J. Solids Structures* 47 (2010) 3115–3120.
- [14] L. Nappa, Variational principles in micromorphic thermoelasticity, *Mech. Res. Commun.* 28 (4) (2001) 405–412.
- [15] J.H. He, A family of variational principles for linear micromorphic elasticity, *Comput. Struct.* 81 (21) (2003) 2079–2085.
- [16] J.H. He, A generalized variational principle in micromorphic thermoelasticity, *Mech. Res. Commun.* 32 (2005) 93–98.
- [17] AC Eringen, Mechanics of micromorphic materials, in: H. Gortler (Ed.), *Proc. 11th Int. Congress of Appl. Mech.*, Springer, New York, 1964.
- [18] J.H. He, Semi-inverse method of establishing generalized variational principles for fluid mechanics with emphasis on turbo machinery aerodynamics, *Int. J. Turbo Jet-Engines* 14 (1) (1997) 23–28.
- [19] J.H. He, Variational theory for one-dimensional longitudinal beam dynamics, *Phys. Lett. A* 352 (4–5) (2006) 276–277.
- [20] J.H. He, E.W.M. Lee, Variational principle for the differential-difference system arising in stratified hydrostatic flows, *Phys. Lett. A* 373 (2009) 1644–1645.
- [21] J.H. He, Variational approach to $(2 + 1)$ -dimensional dispersive long water equations, *Phys. Lett. A* 335 (2005) 182–184.
- [22] C.B. Zheng, B. Liu, Z.J. Wang, et al., Generalized variational principle for electromagnetic field with magnetic monopoles by He's semi-inverse method, *Int. J. Nonlinear Sci. Numer.* 10 (2009) 1369–1372.
- [23] Y. Wu, Variational approach to the generalized Zakharov equations, *Int. J. Nonlinear Sci. Numer.* 10 (2009) 1245–1247.
- [24] S. Pak, Solitary wave solutions for the RLW equation by He's semi-inverse method, *Int. J. Nonlinear Sci. Numer.* 10 (2009) 505–508.
- [25] C.B. Zheng, Z.J. Wang, A Generalized variational principle in special relativity, *Nonlinear Sci. Lett. A* 1 (2010) 245–248.
- [26] Z.J. Wang, D.Z. Zheng, C.B. Zheng, A variational principle for N coupled nonlinear Schrödinger equations, *Int. J. Nonlinear Sci. Numer.* 11 (2010) 401–404.
- [27] B.A. Finlayson, *The Method of Weighted Residuals and Variational Principles*, Acad, New York, 1972.